```
GM Outcome Based Common Standard: Sexual and Reproductive Health
  GM Shared Vision:
 Poor sexual and reproductive health, including the ongoing transmission of HIV, has major impacts on Greater Manchester residents, and despite the progress made, there are still high rates of HIV and STIs in the conurbation. Continuing challenges include the rising rates of some sexually transmitted infe
 outcomes. Almost half all HIV diagnoses in GM are late, which lead to poorer outcomes for the individual and increased risk of onward transmission. Further demands on services are anticipated with the potential introduction of pre-exposure prophylaxis (PrEP) and immediate initiation of anti-retroviral
 • all residents have the knowledge, skills and confidence to make informed choices about their sexual health, reproduction and relationships;
 • sexual and reproductive health services are accessible, sensitive and appropriate for all;
 • improved outcomes in sexual and reproductive health, bringing Greater Manchester to among the best in the country;

    working together to eradicate HIV in a generation

 Our ambition is for a holistic system to ensure good sexual and reproductive health for all Greater Manchester residents with clear pathways, common standards and expectations set within it enabling people to access what they need, at a consistently high quality, when and where they need it. In additing the contract of the contract 
 health and reduce the stigma associated with poor sexual health outcomes. These reforms of the system aim to have the following impacts on the region:
 • GM population will be able to exercise personal choice and self-management regarding sexuality, sexual health and contraception.
 • Significantly reduced prevalence of STIs & HIV in GM, particularly amongst targeted, higher risk communities.
 • Ensure that we are prepared for emerging challenges in sexual health including multidrug resistant gonorrhoea.
 • Improved health and life expectancy for people living with HIV within GM, thus improving the quality of life for people living with HIV and reducing the cost to the sub-region's health and social care system.
 • Maintain open access to sexual and reproductive health services, giving people the choice of where to attend.
 • Agreed standards across the system to ensure that no matter where people gain access to the system, they are able to obtain the right, high quality care.
 • Deliver a more consistent primary care offer, especially for reproductive health.
 Strategic Outcome: Improving the Health of the GM Population and Reducing Health Inequalities across GM
 "I" Statement: "I will live a long and healthy life in Greater Manchester"
                                                                                                                                                                                                                                                             Method of Measuring Impact
                                                                                                      Sexual & Reproductive Health is a strategic consideration within overarching plans for health and social care transformation and
Sexual & Reproductive Health is embedded within Health & Social Care
                                                                                                                                                                                                                                                                           Presence of Sexual & Reproductive Health in plans for Health and Social G
                                                                                                      is embedded within service provision.
 Strategic Outcome: Start Well - Give every GM child the best start in life
 "I" Statement: "I will make sure that every GM child will has the best start in life and will develop well"
                                                                                                                                                                                                                                                             Method of Measuring Impact
                                                                                                                                                                                                                                                                                                                           % of uptake
                                                                                                      All pregnant women are screened for infectious diseases in line with NHS screening guidelines
 Maintain the uptake of syphilis, HIV and Hepatitis B testing in pregnancy
 Strategic Outcome: Live Well - Ensure every GM resident is enabled to fulfil their potential
  "I" Statement: "I will receive an appropriate, non-judgemental service by suitable trained staff"
                                                                                                                                                                                                                                                            Method of Measuring Impact
 Positive patient experience
                                                                                                       Inclusion of questions around sexual & reproductive health in all annual patient surveys (surveys, focus groups)
                                                                                                                                                                                                                                                                                                                         Patient survey
 Patient supported following an HIV diagnosis
 Delivering a responsive service
 "I" Statement: "I will have swift access to the service(s) I need"
                                                                                                                                                                                                                                                            Method of Measuring Impact
 48 hour access to STI treatment and advice for symptomatic patients
                                                                                                                                                                                                                                                                                                                            Clinic data
                                                                                                      100% offer within 48 hours
                                                                                                                                                                                                                                                                                                                    NHS England uptake data
 Improve cervical cancer screening uptake
                                                                                                       80% of women uptake cervial screening
 "I" Statement: "I will be offered choice and support to make an informed decision regarding contraception"
                                                                                                                                                                                                                                                            Method of Measuring Impact
                                                                                                      All under 18s within a locality are encouraged to visit a sexual & reproductive health service or GP before engaging in sexual
                                                                                                                                                                                                                                                                                                               Rate per 1,000 (15-17 year olds)
                                                                                                      activity and are having appropriate levels of contact with these services during adolescence.
                                                                                                      All schools to provide an up-to-date and appropriate age-related RSE programme
Reduction in unwanted pregnancies
                                                                                                      Open access to specialised services for young people up to the age of 19
                                                                                                                                                                                                                                                                            No. of specialist clinic sessions per week for young people available acros
                                                                                                       All young people to have access to school based drop-in sessions
                                                                                                                                                                                                                                                                                            School nurse drop-in sessions available in every secondary
                                                                                                      All women (15-44 years old) are fully informed about and, if clinically appropriate, encouraged to use LARC as their form of
                                                                                                                                                                                                                                                                                                               Rate per 1,000 (15-44 year olds)
Increase in uptake of long acting reversible contraception (LARC)
                                                                                                      For all women having a LARC removed and requiring contraception to have immediate access to an alternative, reliable method
                                                                                                                                                                                                                                                                                                                            Audit (tbc)
   "I" Statement: "I will have access to the testing and treatment I need"
                                                                                                                                                                                                                                                            Method of Measuring Impact
                                                                                                      Routine offer of an HIV test in high prevalence areas and a regular, targeted offer to those in high risk groups
                                                                                                                                                                                                                                                                                                    Number of new diagnoses and % of which are late
 Reduction in new and late diagnosis of HIV
                                                                                                       Evidence of training re Blood Born Viruss for Primary Care every 3 years
                                                                                                                                                                                                                                                                                                Training to GPs/Pharmacies for advice and onward refer
                                                                                                                                                                                                                                                                                              Rate per 100,000 (15-24 year olds) and maintain PN rate of
                                                                                                       Achieve the agreed population level Chlamydia detection rate and meet PN standards
 Improve Chlamydia detection rate
 Reduction in the prevalence of STIs and onward transmission
                                                                                                      Improved digital offer including self-assessment of risk, campaigns
                                                                                                                                                                                                                                                                                                Number of new diagnoses and rate per 100,000 reside
 "I" Statement: "I will be given information and advice about reducing my personal risk of sexual health issues"
                                                                                                                                                                                                                                                            Method of Measuring Impact
 Reduction in abortions and repeat abortions
                                                                                                      LARC offered post-abortion
                                                                                                                                                                                                                                                                                           Rate per 1,000 (15-44 year old women) and % of who are u
 Reduction in repeat STIs
                                                                                                                                                                                                                                                                                                                % reinfected within 12 months
                                                                                                        Provision of personalise risk reduction support and information
       litional relevant guidance for commissioners and providers (i.e NICE Guidance; National Strategy; GM Strategy; Associated GM Common Standards)
NICE Guidance - Sexually transmitted infections and under-18 conceptions:
                                                                                                         https://www.nice.org.uk/guidance/ph3
 prevention [PH3]
NICE Guidance - HIV testing: increasing uptake among people who may have
                                                                                                        HIV testing: increasing uptake among people who may have undiagnosed HIV
 undiagnosed HIV [NG60]

NICE Guidance - Sexually transmitted infections: condom distribution schemes
                                                                                                        https://www.nice.org.uk/guidance/ng68
 NICE Guidance - Harmful sexual behaviour among children and young people [NG55] <a href="https://www.nice.org.uk/guidance/ng55">https://www.nice.org.uk/guidance/ng55</a>
                                                                                                        https://pathways.nice.org.uk/pathways/contraceptive-services-for-under-25s
 NICE Guidance - Contraceptive services for under 25s [PH51]
 NICE Quality Standards - HIV testing: encouraging uptake Quality standard [QS157]
                                                                                                        https://www.nice.org.uk/guidance/qs157
 NICE Quality Standards - Contraception Quality standard [QS129]
                                                                                                        https://www.nice.org.uk/guidance/qs129
 NICE Pathways - Preventing sexually transmitted infections and under-18 conceptions
                                                                                                       https://pathways.nice.org.uk/pathways/preventing-sexually-transmitted-infections-and-under-18-conceptions
                                                                                                        https://pathways.nice.org.uk/pathways/hiv-testing-and-prevention
 NICE Pathways - HIV testing and prevention overview
                                                                                                        https://www.nice.org.uk/guidance/cg30
NICE Guidance - Long Acting Reversible Contraception [CG30]
BHIVA guidelines for the routine investigation and monitoring of adult HIV-1-positive
                                                                                                       http://www.bhiva.org/guidelines.aspx
individuals (2016)
BHIVA guidelines for the treatment of HIV-1-positive adults with antiretroviral therapy

http://www.bhiva.org/HIV-1-treatment-guidelines.aspx
2015 (2016 interim update)
BHIVA guidelines for the management of HIV infection in pregnant women 2012 (2014 <a href="http://www.bhiva.org/pregnancy-guidelines.aspx">http://www.bhiva.org/pregnancy-guidelines.aspx</a>
  interim review)
UK National Guideline for the Use of HIV Post-Exposure Prophylaxis Following Sexual
                                                                                                      http://www.bhiva.org/PEPSE-guidelines.aspx
 Exposure (PEPSE) 2015
 Greater Manchester Sexual & Reprodutive Health Strategy
                                                                                                       http://www.rcgp.org.uk/clinical-and-research/a-to-z-clinical-resources/sexually-transmitted-infections-in-primary-care.aspx
RCGP - Sexually Transmitted Infections in Primary Care
 Faculty of Sexual & Reproductive Health - Contraception Guidelines
                                                                                                         https://www.fsrh.org/standards-and-guidance/current-clinical-guidance/
 Faculty of Sexual & Reproductive Health - Management of SRH Issues Guidelines
                                                                                                          ttps://www.fsrh.org/standards-and-guidance/current-clinical-guidance/management-of-srh-issues/
 NHS Cervical Screening Programme (CSP)
                                                                                                         https://www.gov.uk/topic/population-screening-programmes/cervical
NICE Guidance - Antenatal care for uncomplicated pregnancies [CG62]
                                                                                                        https://www.nice.org.uk/guidance/cg62/ifp/chapter/screening-and-tests
```

nfections, the continuing transmiss al therapy (ART). The vision for G			
ition, the aim to help people be mc			
Care transformation.			
Greater Manchester			
school			
ral			
f 0.6			
ts			
der 25			